

Product datasheet for **MR202089L4V**

Pym1 (NM_001170869) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|--|
| Product Type: | Lentiviral Particles |
| Product Name: | Pym1 (NM_001170869) Mouse Tagged ORF Clone Lentiviral Particle |
| Symbol: | Pym1 |
| Synonyms: | A030010B05Rik; Pym; Wibg |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) |
| Tag: | mGFP |
| ACCN: | NM_001170869 |
| ORF Size: | 609 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(MR202089). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_001170869.1 , NP_001164340.1 |
| RefSeq Size: | 1165 bp |
| RefSeq ORF: | 609 bp |
| Locus ID: | 78428 |
| UniProt ID: | Q8CHP5 |
| Cytogenetics: | 10 D3 |



[View online »](#)

Gene Summary:

Key regulator of the exon junction complex (EJC), a multiprotein complex that associates immediately upstream of the exon-exon junction on mRNAs and serves as a positional landmark for the intron exon structure of genes and directs post-transcriptional processes in the cytoplasm such as mRNA export, nonsense-mediated mRNA decay (NMD) or translation. Acts as an EJC disassembly factor, allowing translation-dependent EJC removal and recycling by disrupting mature EJC from spliced mRNAs. Its association with the 40S ribosomal subunit probably prevents a translation-independent disassembly of the EJC from spliced mRNAs, by restricting its activity to mRNAs that have been translated. Interferes with NMD and enhances translation of spliced mRNAs, probably by antagonizing EJC functions (By similarity). [UniProtKB/Swiss-Prot Function]