

Product datasheet for MR202276

Hmgb2 (BC002050) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Hmgb2 (BC002050) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Hmgb2

Synonyms: C80539; HMG-2; Hmg2

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR202276 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR202276 protein sequence

Red=Cloning site Green=Tags(s)

MGKGDPNKPRGKMSSYAFFVQTCREEHKKKHPDSSVNFAEFSKKCSERWKTMSAKEKSKFEDLAKSDKAR YDREMKNYVPPKGDKKGKKKDPNAPKRPPSAFFLFCSENRPKIKIEHPGLSIGDTAKKLGEMWSEQSAKD KQPYEQKAAKLKEKYEKDIAAYRAKGKSEAGKKGPGRPTGSKKKNEPEDEEEEEEEEEEEEEEEEEEEE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: BC002050

ORF Size: 630 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>BC002050</u>, <u>AAH02050</u>

RefSeq Size: 1076 bp RefSeq ORF: 632 bp



Locus ID: 97165


Cytogenetics: 8 29.9 cM

MW: 24.2 kDa

Gene Summary: Multifunctional protein with various roles in different cellular compartments. May act in a

redox sensitive manner. In the nucleus is an abundant chromatin-associated non-histone protein involved in transcription, chromatin remodeling and V(D)J recombination and probably other processes. Binds DNA with a preference to non-canonical DNA structures such as single-stranded DNA. Can bent DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters by enhancing transcription factor binding and/or bringing distant regulatory sequences into close proximity (By similarity). Involved in V(D)| recombination by acting as a cofactor of the RAG complex: acts by stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS) (PubMed:9184213). Proposed to be involved in the innate immune response to nucleic acids by acting as a cytoplasmic promiscuous immunogenic DNA/RNA sensor which cooperates with subsequent discriminative sensing by specific pattern recognition receptors (PubMed:19890330). In the extracellular compartment acts as a chemokine. Promotes proliferation and migration of endothelial cells implicating AGER/RAGE (By similarity). Has antimicrobial activity in gastrointestinal epithelial tissues (By similarity). Involved in inflammatory response to antigenic stimulus coupled with proinflammatory activity (PubMed:25306442). May play a role in germ cell differentiation (PubMed:11262228). Involved in modulation of neurogenesis probably by regulation of neural stem proliferation (PubMed:24391977). Involved in articular cartilage surface maintenance implicating LEF1 and the Wnt/beta-catenin pathway (PubMed:19805379).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR202276