

## Product datasheet for **MR202338**

### **Bok (NM\_016778) Mouse Tagged ORF Clone**

#### Product data:

Product Type: Expression Plasmids  
 Product Name: Bok (NM\_016778) Mouse Tagged ORF Clone  
 Tag: Myc-DDK  
 Symbol: Bok  
 Synonyms: matador; mtd  
 Mammalian Cell Selection: Neomycin  
 Vector: pCMV6-Entry (PS100001)  
 E. coli Selection: Kanamycin (25 ug/mL)  
 ORF Nucleotide Sequence: >MR202338 ORF sequence  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGAGGTGCTGCGGCGCTCTTCTGTCTTTGCAGCGGAGATCATGGACGCCTTTGATCGCTCGCCACAG  
 ACAAGGAGCTGGTGGCCAGGCTAAGGCACTAGGCCGGGAGTACGTGCACGCGCGCTTTTGCAGCGCGG  
 CCTCTCTGGAGCGCTCCAGAGCGTGCCTCGCTGCCCTGGAGGACGCTTGGCAGAGGTGTGCACAGTG  
 CTGCTGCGCTTGGGAGATGAGCTGGAGCAGATCCGTCCAGCGTATACCGAACGTGGCCCGGCAGCTGC  
 ACATCCCCCTGCAGTCGGAGCCTGTGGTGACGGATGCCTTCTGGCAGTGGCAGGCCACATCTTCTCAGC  
 AGGTATCACATGGGGCAAGGTAGTGTCCCTGTATTCCGTGGCCGCGGGGCTAGCGGTGGACTGTGTCCGG  
 CAGGCTCAGCCTGCCATGGTTTATGCCCTGGTTGACTGCCTGGGGGAATTTGTACGCAAGACCTTGCTA  
 CCTGGCTTCGGAGGCGTGGCGGATGGACGGATGTCTCAAGTGTGTGGTCAAGACAGATCCTGGCTTCCG  
 CTCCCAGTGGCTCGTGGCCACGCTCTGCAGCTTTGGCCGCTTCTGAAGGCAGCATTCTTCTGTTGTTG  
 CCAGAGAGA

**ACGCGT**ACGCGGCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >MR202338 protein sequence  
Red=Cloning site Green=Tags(s)


MEVLRSSVFAAEIMDAFDRSPTDKELVAQAKALGREYVHARLLRAGLSWSAPERASPAPGGRLAEVCTV  
 LLRLGDELEQIRPSVYRNVARQLHIPLQSEPVTDAFLAVAGHIF SAGITWGKVVSLYSVAAGLAVDCVR  
 QAQPAMVHALVDCLGEFVRKTLATWLRRRGGWTDV LKCVVSTDPGFRSHLVATLCSFGRFLKAAFFLLL  
 PER

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



**ACCN:** NM\_016778

**ORF Size:** 642 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_016778.1](#), [NM\\_016778.2](#), [NM\\_016778.3](#), [NP\\_058058.1](#)

**RefSeq Size:** 1432 bp

**RefSeq ORF:** 642 bp

**Locus ID:** 51800


**UniProt ID:** [O35425](#)

**Cytogenetics:** 1 D

**MW:** 23.5 kDa

**Gene Summary:** Apoptosis regulator that functions through different apoptotic signaling pathways (PubMed:23429263, PubMed:26015568, PubMed:26949185, PubMed:27098698, PubMed:9535847). Plays a roles as pro-apoptotic protein that positively regulates intrinsic apoptotic process in a BAX- and BAK1-dependent manner or in a BAX- and BAK1-independent manner (PubMed:23429263, PubMed:26015568, PubMed:26949185). In response to endoplasmic reticulum stress promotes mitochondrial apoptosis through downstream BAX/BAK1 activation and positive regulation of PERK-mediated unfolded protein response (PubMed:26015568). Activates apoptosis independently of heterodimerization with survival-promoting BCL2 and BCL2L1 through induction of mitochondrial outer membrane permeabilization, in a BAX- and BAK1-independent manner, in response to inhibition of ERAD-proteasome degradation system, resulting in cytochrome c release (PubMed:9535847, PubMed:26949185). In response to DNA damage, mediates intrinsic apoptotic process in a TP53-dependent manner. Plays a role in granulosa cell apoptosis by CASP3 activation (By similarity). Plays a roles as anti-apoptotic protein during neuronal apoptotic process, by negatively regulating poly ADP-ribose polymerase-dependent cell death through regulation of neuronal calcium homeostasis and mitochondrial bioenergetics in response to NMDA excitation (PubMed:27098698). In addition to its role in apoptosis, may regulate trophoblast cell proliferation during the early stages of placentar development, by acting on G1/S transition through regulation of CCNE1 expression. May also play a role as an inducer of autophagy by disrupting interaction between MCL1 and BECN1 (By similarity). [UniProtKB/Swiss-Prot Function]

## Product images:



Circular map for MR202338