

Product datasheet for MR202441L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nudt5 (NM_016918) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nudt5 (NM_016918) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nudt5

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_016918

ORF Size: 657 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR202441).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 016918.2</u>, <u>NP 058614.1</u>

 RefSeq Size:
 1590 bp

 RefSeq ORF:
 657 bp

 Locus ID:
 53893

 UniProt ID:
 Q9|KX6

Cytogenetics: 2 A1







Gene Summary:

Enzyme that can either act as an ADP-sugar pyrophosphatase in absence of diphosphate or catalyze the synthesis of ATP in presence of diphosphate (By similarity). In absence of diphosphate, hydrolyzes with similar activities various modified nucleoside diphosphates such as ADP-ribose, ADP-mannose, ADP-glucose, 8-oxo-GDP and 8-oxo-dGDP (PubMed:10722730). Can also hydrolyze other nucleotide sugars with low activity (PubMed:10722730). In presence of diphosphate, mediates the synthesis of ATP in the nucleus by catalyzing the conversion of ADP-ribose to ATP and ribose 5-phosphate (By similarity). Nuclear ATP synthesis takes place when dephosphorylated at Thr-44 (By similarity). Nuclear ATP generation is required for extensive chromatin remodeling events that are energy-consuming (By similarity). Does not play a role in U8 snoRNA decapping activity (PubMed:21070968). Binds U8 snoRNA (PubMed:21070968). [UniProtKB/Swiss-Prot Function]