


Product datasheet for MR202513L3

U2af1l4 (NM_170760) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	U2af1l4 (NM_170760) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	U2af1l4
Synonyms:	AA407033; AF419339; AI451269; AW553050; U2af26
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR202513).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




* The last codon before the Stop codon of the ORF.

ACCN:	NM_170760
ORF Size:	660 bp



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_170760.3 , NP_739566.1
RefSeq Size:	864 bp
RefSeq ORF:	663 bp
Locus ID:	233073
UniProt ID:	Q8BGJ9
Cytogenetics:	7 B1
Gene Summary:	<p>RNA-binding protein that function as a pre-mRNA splicing factor. Plays a critical role in both constitutive and enhancer-dependent splicing by mediating protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. It can functionally substitute for U2AF1 in constitutive splicing and enhancer-dependent splicing. Acts by enhancing the binding of U2AF2 to weak pyrimidine tracts. Also participates in the regulation of alternative pre-mRNA splicing. Activates exon 5 skipping of PTPRC during T-cell activation; an event reversed by GFI1. Binds to RNA at the AG dinucleotide at the 3'-splice site. Shows a preference for AGC or AGA (PubMed:11739736, PubMed:16819553, PubMed:18460468). Alternative splicing of U2AF1L4 may play a role in connecting the circadian rhythm to changing external cues: may provide a circadian buffering system in central and periphery clocks that allows synchronized adaption to clock-resetting stimuli in order to prevent potentially pathogenic desynchronization (PubMed:24837677).[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR202513L3