

Product datasheet for MR202676

Yeats4 (NM_026570) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Yeats4 (NM_026570) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Yeats4

Synonyms: 4930573H17Rik; B230215M10Rik; GAS41; NuBl-1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>MR202676 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGTTCAAGAGAATGCCGGAATTTGGACCTGACTCCGGCGGGAGAGTGAAGGGGGTTACCATCGTTAAGC
CAATCGTTTATGGCAATGTTGCCAGGTACTTTGGAAAGAAGAAGAAGAAGAAGAAGAAGACGGCACACTCACCAGTG
GACTGTGTACGTGAAGCCCTACAGAAACGAGGATATGTCAGCATATGTGAAGAAGACCAGTTTAAATTA
CACGAAAGCTACGGCAATCCTCTAAGAGTCGTCACCAAGCCTCCATATGAAATCACAGAAACAGGATGGG
GTGAATTTGAAATCATCATCAAGATATTTTTCATTGACCCCAACGAGAGACCTGTGACTCTGTACCACTT
ACTGAAGCTCTTCCAGTCTGACACCAATGCCATGCTGGGGAAGAAAACAGTGGTTTCAGAGTTCTATGAC
GAAATGATATTTCAAGACCCAACGGCAATGATGCAGCAGCTGCTTACGACGTCTCGCCAGCTGACATTGG
GAGCCTATAAGCACGAAACAGAGTTTGCAGAACTTGAAGTGAAAACCAGAGAAAAAATTAGAAGCTGCCAA
GAAAAAAAACAAGCTTTGAGATTGCGGAGCTTAAGGAGAGATTAAAACCAGTCGTGAAACTATAAACTGT
TTAAAAAAATGAAATCAGGAAGCTTGAAGAAGACGATCAGACGAAAGACATT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >MR202676 protein sequence

Red=Cloning site Green=Tags(s)


MFKRMAEFGPDSGGRVKGVTIVKPIVYGNVARYFGKKREEDGHTHQWTVYVKPYRNEDMSAYVKKIQFKL HESYGNPLRVVTKPPYEITETGWGEFEIIIKIFFIDPNERPVTLYHLLKLFQSDTNAMLGKKTVVSEFYD EMIFQDPTAMMQQLLTTSRQLTLGAYKHETEFAELEVKTREKLEAAKKKTSFEIAELKERLKASRETINC LKNEIRKLEEDDQTKDI


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_026570

ORF Size: 681 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 026570.4</u>

 RefSeq Size:
 1418 bp

 RefSeq ORF:
 684 bp

 Locus ID:
 64050

 UniProt ID:
 Q9CR11

 Cytogenetics:
 10 D2

 MW:
 26.5 kDa

Gene Summary: Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in


transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote

interaction of the modified histones with other proteins which positively regulate

transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage (By similarity).

[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR202676

