

Product datasheet for MR202750L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Asah3l (BC059819) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Asah3l (BC059819) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Asah3

Synonyms: CRG-L1, Acer2, maCER2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 BC059819

 ORF Size:
 687 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR202750).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally accurring varieties (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: BC059819, AAH59819

RefSeq Size:4078 bpRefSeq ORF:689 bpLocus ID:230379

Cytogenetics: 4 C4







Gene Summary:

Golgi ceramidase that catalyzes the hydrolysis of ceramides into sphingoid bases like sphingosine and free fatty acids at alkaline pH (PubMed:29401619). Ceramides, sphingosine, and its phosphorylated form sphingosine-1-phosphate are bioactive lipids that mediate cellular signaling pathways regulating several biological processes including cell proliferation, apoptosis and differentiation (PubMed:29401619). Has a better catalytic efficiency towards unsaturated long-chain ceramides, including C18:1-, C20:1- and C24:1-ceramides (By similarity) (PubMed:29401619). Saturated long-chain ceramides and unsaturated very longchain ceramides are also good substrates, whereas saturated very long-chain ceramides and short-chain ceramides are poor substrates. Also hydrolyzes dihydroceramides to produce dihydrosphingosine (By similarity). It is the ceramidase that controls the levels of circulating sphingosine-1-phosphate and dihydrosphingosine-1-phosphate in plasma through their production by hematopoietic cells (PubMed:29401619). Regulates cell proliferation, autophagy and apoptosis by the production of sphingosine and sphingosine-1-phosphate. As part of a p53/TP53-dependent pathway, promotes for instance autophagy and apoptosis in response to DNA damage. Through the production of sphingosine, may also regulate the function of the Golgi complex and regulate the glycosylation of proteins (By similarity). [UniProtKB/Swiss-Prot Function]