

## Product datasheet for **MR202793L3V**

### **Nqo2 (NM\_001163242) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Nqo2 (NM_001163242) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Nqo2
Synonyms:	NMO2; Nmor2; Ox2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001163242
ORF Size:	696 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR202793).
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001163242.1</a> , <a href="#">NP_001156714.1</a>
RefSeq Size:	3821 bp
RefSeq ORF:	696 bp



[View online »](#)

Locus ID: 18105

UniProt ID: [Q9JI75](#)

Cytogenetics: 13 14.01 cM

**Gene Summary:** The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.[UniProtKB/Swiss-Prot Function]