

## Product datasheet for **MR202802L1V**

### **Klrk1 (NM\_033078) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | Klrk1 (NM_033078) Mouse Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | Klrk1  |
| Synonyms:                 | D6H12S2489E; NKG2-D; Nkg2d   |
| Mammalian Cell Selection: | None   |
| Vector:                   | pLenti-C-Myc-DDK (PS100064)  |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_033078  |
| ORF Size:                 | 699 bp   |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR202802).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_033078.3</a>  |
| RefSeq Size:              | 3272 bp  |
| RefSeq ORF:               | 699 bp   |
| Locus ID:                 | 27007  |
| UniProt ID:               | <a href="#">O54709</a>   |
| Cytogenetics:             | 6 63.44 cM   |



[View online »](#)

**Gene Summary:**

Function as an activating and costimulatory receptor involved in immunosurveillance upon binding to various cellular stress-inducible ligands displayed at the surface of autologous tumor cells and virus-infected cells. Provides both stimulatory and costimulatory innate immune responses on activated killer (NK) cells, leading to cytotoxic activity. Acts as a costimulatory receptor for T-cell receptor (TCR) in CD8(+) T-cell-mediated adaptive immune responses by amplifying T-cell activation. Stimulates perforin-mediated elimination of ligand-expressing tumor cells. Signaling involves calcium influx, culminating in the expression of TNF-alpha. Participates in NK cell-mediated bone marrow graft rejection. May play a regulatory role in differentiation and survival of NK cells. Binds to ligands belonging to various subfamilies of MHC class I-related glycoproteins including RAET1A, RAET1B, RAET1C, RAET1D, RAET1E, H60 and MULT1.[UniProtKB/Swiss-Prot Function]