

## Product datasheet for **MR202922L4V**

### **Cd63 (NM\_001042580) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Cd63 (NM_001042580) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cd63
Synonyms:	C75951; ME491; Tspan30
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001042580
ORF Size:	717 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR202922).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001042580.1</a> , <a href="#">NP_001036045.1</a>
RefSeq Size:	959 bp
RefSeq ORF:	717 bp
Locus ID:	12512
UniProt ID:	<a href="#">P41731</a>
Cytogenetics:	10 77.19 cM



[View online »](#)

**Gene Summary:**

Functions as cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2. Plays a role in VEGFA signaling via its role in regulating the internalization of KDR/VEGFR2. Plays a role in intracellular vesicular transport processes, and is required for normal trafficking of the PMEL luminal domain that is essential for the development and maturation of melanocytes. Plays a role in the adhesion of leukocytes onto endothelial cells via its role in the regulation of SELP trafficking. May play a role in mast cell degranulation in response to Ms4a2/FcεRI stimulation, but not in mast cell degranulation in response to other stimuli.[UniProtKB/Swiss-Prot Function]