

Product datasheet for MR203320L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Zfand2b (NM_001159905) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Zfand2b (NM 001159905) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Zfand2b

Synonyms: 1110060O18Rik; C81256

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001159905

ORF Size: 774 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR203320).

OTI Disclaimer:

Cytogenetics:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001159905.1</u>, <u>NP 001153377.1</u>

1 C4

 RefSeq Size:
 1206 bp

 RefSeq ORF:
 774 bp

 Locus ID:
 68818

 UniProt ID:
 Q91X58





Gene Summary:

Plays a role in protein homeostasis by regulating both the translocation and the ubiquitin-mediated proteasomal degradation of nascent proteins at the endoplasmic reticulum (PubMed:24160817, PubMed:26337389, PubMed:26692333). It is involved in the regulation of signal-mediated translocation of proteins into the endoplasmic reticulum (PubMed:24160817). It also plays a role in the ubiquitin-mediated proteasomal degradation of proteins for which signal-mediated translocation to the endoplasmic reticulum has failed (PubMed:18467495, PubMed:26337389). May therefore function in the endoplasmic reticulum stress-induced pre-emptive quality control, a mechanism that selectively attenuates the translocation of newly synthesized proteins into the endoplasmic reticulum and reroutes them to the cytosol for proteasomal degradation (PubMed:24160817, PubMed:26337389). By controlling the steady-state expression of the IGF1R receptor, indirectly regulates the insulin-like growth factor receptor signaling pathway (PubMed:26692333).[UniProtKB/Swiss-Prot Function]