

Product datasheet for MR203331

Eif4e2 (BC049077) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Eif4e2 (BC049077) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Eif4e2
Synonyms:	2700069E09Rik; AI036339; AV129531; D0H0S6743E; Eif4el3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR203331 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAACAACAAGTTCGACGCCTTAAAAGATGATGACAGTGGAGACCATGATCAGAATGAAGAAAACAGCA
CACAGAAAGATGGTGAGAAGGAAAAACAGACCGAGACAAGAGCCAGAGCAGTGGCAAGAGGAAGGCTGT
TGTCCTGGACCAGCAGAGCATCCCTGCAGTACAACCTACACCTTTTGGTACTCGAGGAGAACCCCTGGC
CGTCCCACCAGCTCGCAGAGCTATGAGCAGAACATCAAGCAGATTGGCACCTTTGCCTCTGTGGAGCAGT
TCTGGAAGTTTTACAGCCACATGGTACGTCCTGGGGACCTGACAGGCCACAGTGACTTTTCATCTCTCAA
AGAAGGGATTAACCTATGTGGGAGGATGATGCAAATAAAAAATGGGGCAAGTGGATCATTTCGACTCCGG
AAGGGCTTAGCTTCCCGCTGCTGGGAGAATCTCATCTGGCTATGCTCGGGGAGCAATTCATGGTTGGGG
AGGAGATCTGCGGGCTGTGGTCTCTGTCCGCTTTCAGGAGGACATTATTTCTATATGGAATAAGACTGC
CAGCGACCAAGCAACTACAGCCGAATCCGGGATACTTTCGGCGCGTGCTTAACCTACCTCCCAACACC
ATTATGGAATACAAAACCTACACCGACAGCATTGCACCTCAGGGAACCATGAGTTCCAAACTATGTCTCA
TGGTTAACCCCTTAAAGGTCACCAGACCTCATGCTTCTACGTTCCCGCTTGGCTCCGTGCACCAACAGAT
C

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR203331 protein sequence
Red=Cloning site Green=Tags(s)

MNNKFDALKDDDSGDHDQNEENSTQKDGEKEKTDRDKSQSSGKRKAVVPGPAEHPLQYNYTFWYSRRTPG
 RPTSSQSYEQNIKQIGTFASVEQFWKFYSHMVRPGDLTGHSDFHFLKEGIKPMWEDDANKNGGKWIIRLR
 KGLASRCWENLILAMLGEQFMVGEEICGAVVSVRFQEDIISIWNKTASDQATTARIRDTLRRVNLNLPNT
 IMEYKHTHTDSIAPQGTMSKLCMLMVNPLKVTRPHASTFPLGSVHQQI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: BC049077

ORF Size: 771 bp


OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>BC049077</u> , <u>AAH49077</u>
RefSeq Size:	2855 bp
RefSeq ORF:	773 bp
Locus ID:	26987
Cytogenetics:	1 C5
MW:	29.3 kDa
Gene Summary:	Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation (PubMed:15153109). Acts as a repressor of translation initiation (By similarity). In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block assembly of eIF4F at the cap (PubMed:15153109). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR203331