

Product datasheet for MR203545L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Gpm6a (BC023461) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Gpm6a (BC023461) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Gpm6a

Synonyms: MGC38999, M6A

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: BC023461 **ORF Size:** 801 bp

ORF Nucleotide

Sequence:
OTI Disclaimer:

Cytogenetics:

The ORF insert of this clone is exactly the same as(MR203545).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: BC023461, AAH23461

8 B1.3

RefSeq Size:2780 bpRefSeq ORF:803 bpLocus ID:234267





Gene Summary:

Involved in neuronal differentiation, including differentiation and migration of neuronal stem cells. Plays a role in neuronal plasticity and is involved in neurite and filopodia outgrowth, filopodia motility and probably synapse formation. Gpm6a-induced filopodia formation involves mitogen-activated protein kinase (MAPK) and Src signaling pathways. Conflictingly, PubMed:22162747 reports that induced cellular protrusions are simple membrane-wrapped tubules without actin or tubulin-based cytoskeletons and with Gpm6a gliding along membrane edges indicative for a function in actin-independent membrane deformation. May be involved in neuronal NGF-dependent Ca(2+) influx. May be involved in regulation of endocytosis and intracellular trafficking of G-protein-coupled receptors (GPCRs); enhances internalization and recycling of mu-type opioid receptor.[UniProtKB/Swiss-Prot Function]