

Product datasheet for **MR203549L3V**

Med8 (NM_020000) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | Lentiviral Particles |
| Product Name: | Med8 (NM_020000) Mouse Tagged ORF Clone Lentiviral Particle |
| Symbol: | Med8 |
| Synonyms: | 2210021A15Rik; AB041805; ARC32 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| Tag: | Myc-DDK |
| ACCN: | NM_020000 |
| ORF Size: | 807 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(MR203549). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_020000.2 , NP_064384.2 |
| RefSeq Size: | 1887 bp |
| RefSeq ORF: | 807 bp |
| Locus ID: | 80509 |
| UniProt ID: | Q9D7W5 |
| Cytogenetics: | 4 |



[View online »](#)

Gene Summary:

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. May play a role as a target recruitment subunit in E3 ubiquitin-protein ligase complexes and thus in ubiquitination and subsequent proteasomal degradation of target proteins (By similarity).[UniProtKB/Swiss-Prot Function]