

Product datasheet for MR203694L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Psmb8 (NM_010724) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Psmb8 (NM 010724) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Psmb8

Synonyms: Lmp-7; Lmp7

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 010724

ORF Size: 831 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR203694).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally accurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 010724.1, NP 034854.1

RefSeq Size: 1223 bp
RefSeq ORF: 831 bp
Locus ID: 16913
UniProt ID: P28063

Cytogenetics: 17 17.98 cM







Gene Summary:

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. This subunit is involved in antigen processing to generate class I binding peptides. May participate in the inflammatory response pathway. Required for adipocyte differentiation (PubMed:21881205, PubMed:22341445, PubMed:8066463). May be involved in the generation of spliced peptides resulting from the ligation of two separate proteasomal cleavage products that are not contiguous in the parental protein (By similarity).[UniProtKB/Swiss-Prot Function]