

Product datasheet for MR203949L4V

Ccnd2 (NM_009829) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ccnd2 (NM_009829) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Ccnd2
Synonyms:	2600016F06Rik; Al256817; BF642806; C86853; cD2; Vin-1; Vin1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_009829
ORF Size:	870 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR203949).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 009829.3, NP 033959.1</u>
RefSeq Size:	5772 bp
RefSeq ORF:	870 bp
Locus ID:	12444
UniProt ID:	<u>P30280</u>
Cytogenetics:	6 61.92 cM



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



Gene Summary:Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits
members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle
during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor
E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are
responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early
G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and
antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-
dependent manner and repressing its transcriptional activity. Component of the ternary
complex, cyclin D2/CDK4/CDKN1B, required for nuclear translocation and activity of the
cyclin D-CDK4 complex (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US