

Product datasheet for MR203964L4

1110004E09Rik (NM_026502) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	1110004E09Rik (NM_026502) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	1110004E09Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR203964).
Restriction Sites:	Sgfl-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_026502
ORF Size:	873 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_026502.1 , NP_080778.1
RefSeq Size:	1163 bp
RefSeq ORF:	873 bp
Locus ID:	68001
UniProt ID:	Q8BL95
Cytogenetics:	16 C3.3
Gene Summary:	Plays a role in motile cilium function, possibly by acting on outer dynein arm assembly. Seems to be important for initiation rather than maintenance of cilium motility. Required for correct positioning of cilia at the apical cell surface, suggesting an additional role in the planar cell polarity (PCP) pathway. May suppress canonical Wnt signaling activity.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR203964L4