

## Product datasheet for MR204015

### Hnrnpc (NM\_001170984) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Hnrnpc (NM_001170984) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Hnrnpc
Synonyms:	AL022939; D14Wsu171e; hnrnp-C; hnRNPC1; hnRNPC2; Hnrpc; Hnrpc1; Hnrpc2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR204015 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCTAGCAATGTTACCAACAAGACAGATCCTCGGTCCATGAATCCCCTGTATTCATTGGGAATCTCA  
ACACTCTGGTGGTCAAGAAGTCTGATGTGGAGGCCATCTTTCAAAGTATGGCAAATTGTGGGCCGCTC  
TGTGCATAAAGGCTTTGCCTTTGTCCAGTATGTTAATGAAAGAAATGCCCGAGCTGCTGTAGCTGGCGAG  
GATGGCAGAATGATTGCTGGCCAGGTTTTAGATATTAACCTGGCTGCAGAGCCAAAAGTGAACCGAGGAA  
AAGCAGGTGTGAAACGATCTGCAGCGGAGATGTACGGTTCCTCATTGACTTGGACTATGACTTTCAACG  
GGATTATTATGACAGGATGTACAGTTACCCAGCGCGGGTTCCTCCTCCTCCTATTGCTCGAGCTGTG  
GTGCCTTCCAAACGTCAGCGTGTTCAGGGAACACCTCACGAAGGGGCAAAGTGGATTCAATTCGAAGA  
GTGGACAAAGGGGATCTTCTTCCAAGTCTGAAAATTGAAAGGTGATGACCTTCAGGCCATTAATAAGGA  
GCTGACTCAGATAAAACAAAAGTGGATTCTTCTTCTGAAAGCCTGGAAAAATTGAAAAAGAACAAAGC  
AAGCAAGCAGAGATGAAGAATGAAAAGTCTGAAGAAGAGCAGAGCAGCGCCTCTGTGAAGAAAGATGAGA  
CTAATGTGAAGATGGAGTCTGAGGCAGGTGCAGATGACTCTGCTGAGGAGGGTGACCTGCTGGATGATGA  
CGATAATGAAGATCGGGGGATGACCAGCTGGAGTTGAAGGATGATGAAAAAGAGCCTGAGGAAGGAGAA  
GACGACAGAGACAGCCCAATGGGGAGGATGACTCT

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >MR204015 protein sequence  
 Red=Cloning site Green=Tags(s)

MASNVTNKTDPRSMNSRVFIGNLNLTLVVKKSDVEAIFSKYGKIVGRSVHKGFVQYVNERNARA AVAGE  
 DGRMIAGQVLDINLAAEPKVNVRGKAGVKRSAAEMYGSSFDLDYDFQRDYDRMYSYPARVPPPPPIARAV  
 VPSKRQRVSGNTSRRGKSGFNKSGQRGSSSKSGKLGDDLQA IKKELTQIKQKVDLSLESLEKIEKEQS  
 KQAEMKNEKSEEEQSSASVKKDETNVKMESEAGADDSAEEDGLDDDDNEDRGDDQLELKDDEKEPEEGE  
 DDRDSANGEDDS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF

**ACCN:** NM\_001170984

**ORF Size:** 879 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001170984.1](#), [NP\\_001164455.1](#)

**RefSeq Size:** 2781 bp

**RefSeq ORF:** 879 bp

**Locus ID:** 15381


**UniProt ID:** [Q9Z204](#)

**Cytogenetics:** 14 26.79 cM

**MW:** 32.3 kDa

**Gene Summary:** Binds pre-mRNA and nucleates the assembly of 40S hnRNP particles. Interacts with poly-U tracts in the 3' UTR or 5'-UTR of mRNA and modulates the stability and the level of translation of bound mRNA molecules. Single HNRNPC tetramers bind 230-240 nucleotides. Trimers of HNRNPC tetramers bind 700 nucleotides. May play a role in the early steps of spliceosome assembly and pre-mRNA splicing. N6-methyladenosine (m6A) has been shown to alter the local structure in mRNAs and long non-coding RNAs (lncRNAs) via a mechanism named 'm(6)A-switch', facilitating binding of HNRNPC, leading to regulation of mRNA splicing. [UniProtKB/Swiss-Prot Function]

### Product images:



Circular map for MR204015