

## Product datasheet for **MR204036L2V**

### Fcgr2b (NM\_010187) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Fcgr2b (NM_010187) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Fcgr2b
Synonyms:	AI528646; CD32; F630109E10Rik; Fcgr2; Fcgr2a; FcgRII; Fcr-2; Fcr-3; fcRII; Fc[g]RII; Ly-17
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_010187
ORF Size:	882 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR204036).
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_010187.2</a> , <a href="#">NP_034317.1</a>
RefSeq Size:	1415 bp
RefSeq ORF:	882 bp



[View online »](#)

Locus ID: 14130

UniProt ID: [P08101](#)

Cytogenetics: 1 78.02 cM

**Gene Summary:** Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of antigen-antibody complexes from the circulation and modulation of antibody production by B-cells. Isoform IIB1 and isoform IIB1' form caps but fail to mediate endocytosis or phagocytosis. Isoform IIB2 can mediate the endocytosis of soluble immune complexes via clathrin-coated pits. Isoform IIB1 and isoform IIB2 can down-regulate B-cell, T-cell, and mast cell activation when coaggregated to B-cell receptors for AG (BCR), T-cell receptors for AG (TCR), and Fc receptors, respectively.[UniProtKB/Swiss-Prot Function]