

Product datasheet for MR204085L4

Cfdp1 (NM_011801) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cfdp1 (NM_011801) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Cfdp1
Synonyms:	AA408409; Bcnt; Bucentaur; Cfdp; cp27
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR204085).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_011801
ORF Size:	888 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_011801.1
RefSeq Size:	1215 bp
RefSeq ORF:	888 bp
Locus ID:	23837
UniProt ID:	O88271
Cytogenetics:	8 E1
Gene Summary:	May play a role during embryogenesis. May modulate tooth organogenesis since alterations of this protein function affect tooth organs size as well as individual cell fate and survival. In embryonic cells, blockage of the function results in increased number of apoptotic cells, reduced proliferation, alterations in cell shape and fibronectin matrix synthesis. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR204085L4