

## Product datasheet for MR204098L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Plscr3 (NM\_023564) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Plscr3 (NM\_023564) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Plscr3

**Synonyms:** 2210403O21Rik; 2610037N06Rik; ESTM3; X83310

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_023564

ORF Size: 888 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR204098).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 023564.2

 RefSeq Size:
 2785 bp

 RefSeq ORF:
 891 bp

 Locus ID:
 70310

 UniProt ID:
 Q9||Z9

Cytogenetics: 11 42.9 cM







## **Gene Summary:**

May mediate accelerated ATP-independent bidirectional transbilayer migration of phospholipids upon binding calcium ions that results in a loss of phospholipid asymmetry in the plasma membrane. May play a central role in the initiation of fibrin clot formation, in the activation of mast cells and in the recognition of apoptotic and injured cells by the reticuloendothelial system. Seems to play a role in apoptosis, through translocation of cardiolipin from the inner to the outer mitochondrial membrane which promotes BID recruitment and enhances tBid-induced mitochondrial damages.[UniProtKB/Swiss-Prot Function]