

## Product datasheet for MR204260L4

### Crk (NM\_133656) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Crk (NM_133656) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Crk
Synonyms:	c-Crk; Cr; Crk-I; Crk-II; Crk-III; Crk3; CrkIII; Crko; p38
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR204260).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_133656
ORF Size:	915 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_133656.2</a>
<b>RefSeq Size:</b>	6005 bp
<b>RefSeq ORF:</b>	915 bp
<b>Locus ID:</b>	12928
<b>UniProt ID:</b>	<a href="#">Q64010</a>
<b>Cytogenetics:</b>	11 45.92 cM
<b>Gene Summary:</b>	This gene is part of a family of adapter proteins that mediate formation of signal transduction complexes in response to extracellular stimuli, such as growth and differentiation factors. Protein-protein interactions occur through the SH2 domain, which binds phosphorylated tyrosine residues, and the SH3 domain, which binds proline-rich peptide motifs. These interactions promote recruitment and activation of effector proteins to regulate cell migration, adhesion, and proliferation. In mouse this protein is essential for embryonic development. Alternatively spliced transcripts encoding different isoforms with distinct biological activity have been described. [provided by RefSeq, Mar 2013]

Product images:



Circular map for MR204260L4