

# **Product datasheet for MR204260L4V**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Crk (NM\_133656) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Crk (NM\_133656) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Crk

Synonyms: c-Crk; Cr; Crk-I; Crk-II; Crk-III; Crk3; CrkIII; Crko; p38

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_133656

ORF Size: 915 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR204260).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 133656.2

 RefSeq Size:
 6005 bp

 RefSeq ORF:
 915 bp

 Locus ID:
 12928

 UniProt ID:
 Q64010

Cytogenetics: 11 45.92 cM







### **Gene Summary:**

This gene is part of a family of adapter proteins that mediate formation of signal transduction complexes in response to extracellular stimuli, such as growth and differentiation factors. Protein-protein interactions occur through the SH2 domain, which binds phosphorylated tyrosine residues, and the SH3 domain, which binds proline-rich peptide motifs. These interactions promote recruitment and activation of effector proteins to regulate cell migration, adhesion, and proliferation. In mouse this protein is essential for embryonic development. Alternatively spliced transcripts encoding different isoforms with distinct biological activity have been described. [provided by RefSeq, Mar 2013]