

## Product datasheet for MR204283L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Ubb (NM\_011664) Mouse Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Ubb (NM\_011664) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ubb

Synonyms: AL033289; Rps27a; Uba52; Ubb2; Ubc

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 011664

ORF Size: 918 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR204283).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 011664.2</u>, <u>NP 035794.1</u>

RefSeq Size: 1165 bp
RefSeq ORF: 918 bp
Locus ID: 22187
UniProt ID: POCG49

Cytogenetics: 11 38.46 cM







## **Gene Summary:**

This gene encodes ubiquitin, one of the most conserved proteins known. Ubiquitin has a major role in targeting cellular proteins for degradation by the 26S proteosome. It is also involved in the maintenance of chromatin structure, the regulation of gene expression, and the stress response. Ubiquitin is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin moiety fused to an unrelated protein. This gene consists of four direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. Pseudogenes of this gene are located on chromosomes 3 and 14. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]