

## Product datasheet for MR204456L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Rnf126 (NM\_144528) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Rnf126 (NM\_144528) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Rnf126

**Synonyms:** 2610010O19Rik

Mammalian Cell

Puro

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_144528

ORF Size: 942 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR204456).

Sequence:

Cytogenetics:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 144528.1

10 C1

 RefSeq Size:
 1639 bp

 RefSeq ORF:
 942 bp

 Locus ID:
 70294

 UniProt ID:
 Q91YL2





### **Gene Summary:**

E3 ubiquitin-protein ligase that mediates ubiquitination oF target proteins (By similarity). Depending on the associated E2 ligase, mediates 'Lys-48'- and 'Lys-63'-linked polyubiquitination of substrates (PubMed:23418353). Part of a BAG6-dependent quality control process ensuring that proteins of the secretory pathway that are mislocalized to the cytosol are degraded by the proteasome (By similarity). Probably acts by providing the ubiquitin ligase activity associated with the BAG6 complex and be responsible for ubiquitination of the hydrophobic mislocalized proteins and their targeting to the proteasome (By similarity). May also play a role in the endosomal recycling of IGF2R, the cation-independent mannose-6-phosphate receptor (By similarity). May play a role in the endosomal sorting and degradation of several membrane receptors including EGFR, FLT3, MET and CXCR4, by mediating their ubiquitination (PubMed:23418353). By ubiquitinating CDKN1A/p21 and targeting it for degradation, may also promote cell proliferation (By similarity). May monoubiquitinate AICDA (By similarity).[UniProtKB/Swiss-Prot Function]