

Product datasheet for MR204539L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Polr3f (NM_029763) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Polr3f (NM_029763) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Polr3f

Synonyms: 2810411G20Rik; 3010019O03Rik; 3110032A07Rik; RPC6; RPC39

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_029763

ORF Size: 951 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR204539).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 029763.3</u>

RefSeq Size: 4160 bp
RefSeq ORF: 951 bp
Locus ID: 70408
UniProt ID: Q921X6
Cytogenetics: 2 G1







Gene Summary:

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Specific peripheric component of RNA polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. May direct RNA Pol III binding to the TFIIIB-DNA complex. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts induce type I interferon and NF-Kappa-B through the RIG-I pathway (By similarity). Preferentially binds double-stranded DNA (dsDNA) (By similarity).[UniProtKB/Swiss-Prot Function]