

## Product datasheet for MR204670L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Ppp1cc (BC010613) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Ppp1cc (BC010613) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ppp1cc

Synonyms: dis2m1, PP1

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 BC010613

 ORF Size:
 969 bp

**ORF Nucleotide** 

NDE Nucleatide The Of

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR204670).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** BC010613, AAH10613

RefSeq Size:2166 bpRefSeq ORF:971 bpLocus ID:19047

Cytogenetics: 5 F







### **Gene Summary:**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Dephosphorylates RPS6KB1. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E.[UniProtKB/Swiss-Prot Function]