

Product datasheet for MR204893L4

Gpr18 (NM_182806) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Gpr18 (NM_182806) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Gpr18
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR204893).
Restriction Sites:	Sgfl-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_182806
ORF Size:	996 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_182806.1 , NP_877958.1
RefSeq Size:	1334 bp
RefSeq ORF:	996 bp
Locus ID:	110168
UniProt ID:	Q8K1Z6
Cytogenetics:	14 65.86 cM
Gene Summary:	Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (By similarity). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:23104136). Can also be activated by plant-derived and synthetic cannabinoid agonists (By similarity). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (By similarity). May contribute to regulation of the immune system (By similarity). Is required for normal homeostasis of CD8+ subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs) in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (PubMed:25348153, PubMed:26197390). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (PubMed:23461720, PubMed:27893106). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR204893L4