

Product datasheet for **MR204911L4V**

Six3 (NM_011381) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Six3 (NM_011381) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Six3
Synonyms:	E130112M24Rik; Six3a; Six3alpha; Six3b; Six3beta
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_011381
ORF Size:	999 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR204911).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_011381.4 , NP_035511.2
RefSeq Size:	3680 bp
RefSeq ORF:	1002 bp
Locus ID:	20473
UniProt ID:	Q62233
Cytogenetics:	17 55.42 cM


[View online »](#)

Gene Summary:

Transcriptional regulator which can act as both a transcriptional repressor and activator by binding a ATTA homeodomain core recognition sequence on these target genes. During forebrain development represses WNT1 expression allowing zona limitans intrathalamica formation and thereby ensuring proper antero-posterior patterning of the diencephalon and formation of the rostral diencephalon (PubMed:18094027). Acts as a direct upstream activator of SHH expression in the rostral diencephalon ventral midline and that in turn SHH maintains its expression (PubMed:18775421). In addition, Six3 activity is required for the formation of the telencephalon. During postnatal stages of brain development is necessary for ependymal cell maturation by promoting the maturation of radial glia into ependymal cells through regulation of neuroblast proliferation and migration (PubMed:22071110). Acts on the proliferation and differentiation of neural progenitor cells through activating transcription of CCND1 AND CCND2 (PubMed:17576749). During early lens formation plays a role in lens induction and specification by activating directly PAX6 in the presumptive lens ectoderm (PubMed:17066077). In turn PAX6 activates SIX3 resulting in activation of PDGFRA and CCND1 promoting cell proliferation (PubMed:12072567). Also is required for the neuroretina development by directly suppressing WNT8B expression in the anterior neural plate territory (PubMed:20890044). Its action during retina development and lens morphogenesis is TLE5 and TLE4-dependent manner. Furthermore, during eye development regulates several genes expression. Before and during early lens development represses the CRYGF promoter by binding a SIX repressor element (PubMed:11139622). Directly activates RHO transcription, or cooperates with CRX or NRL (PubMed:17666527). Six3 functions also in the formation of the proximodistal axis of the optic cup (PubMed:12163408), and promotes the formation of optic vesicles-like structures (PubMed:11458394). During pituitary development, acts in parallel or alternatively with HESX1 to control cell proliferation through Wnt/beta-catenin pathway (PubMed:18694563). Plays a role in eye development by suppressing WNT1 expression and in dorsal-ventral patterning by repressing BMP signaling pathway (By similarity).[UniProtKB/Swiss-Prot Function]