

Product datasheet for MR204929

Babam1 (NM_026636) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Babam1 (NM_026636) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Babam1
Synonyms:	5430437P03Rik; Merit40
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR204929 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAGGTGGCAGAGGCCAACAGCCCCACGGAGGAGGAGGAAGAGGAGGAAGAAGAAGGAGAGGAGCCGA
TATCAGAGCCCAGGCCTCACACTCGATCCAATCCAGAGGGGGCTGAGGACCGGCCCTGGGAGCTCAAGC
CAATGTGGGCAGCCGAGCGAGGGCGAGGGTGAGGCAGCCACTGCAGATGGCGGGCAGCCAGTGTCCCA
GGAGCCGGGCCTAAGCCCTGGCAAGTACCAGCATCAGCATCTGAGGTCCAGATTCGAACACCAAGGGTCA
ACTGTCCAGAGAAGGTGATCATCTGTCTGGATCTTTCGGAGGAGATGTCTGTGCCAAAGCTGGAGTCCTT
TAATGGCTCTAGGACAAACGCCCTGAATGTGTCTCAGAAGATGGTTGAGATGTTTGTTCGCACAAAGCAC
AAGATTGACAAGAGCCACGAATTTGCACTGGTTGTAGTGAATGACGACTCCGCCTGGCTGTCTGGCCTGA
CCTCTGACCCACGTGAGCTCTGCAGTGCCTGTATGACCTGGAGACGGCGTCTGTCTCCACCTTCAATTT
GGAAGGCCTCTTCAGCCTCATCCAGCAGAAGACCGAGCTGCCAGTCACAGAGAATGTGCAAACCATCCCA
CCCCCTATGTTGTGCGCACCATCCTGGTCTACAGCCGCCACCTGCCAGCCCCAGTTTTCCCTGACTG
AGCCCATGAAGAAGATGTTCCAGTGTCCCTACTTCTTCTCGACATCGTTTATATCCACAATGGCACTGA
GGAGAAGGAAGAGGATATGAGCTGGAAGGACATGTTTGCCTTCATGGGTAGCCTGGACACCAAGGGCGCC
AGCTACAAGTATGAGGTAGCACTTGTCTGGTCTGCCCTGGAGCTGCACAATTGCATGGCCAAGTTGCTGG
CCACCCGCTGCAAAGACCTGCCAGACCCACTCGAGCTACAGCCTGCTGGAGGAGGACGAGGAGGCTGG
AGAAGAGGAAGCCACTGTG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR204929 protein sequence
Red=Cloning site Green=Tags(s)

```
MEVAEANSPTEEEEEEEEEEGEEPISEPRPHTRS NPEGAEDRALGAQANVGSRSEGEAATADGGAASVP
GAGPKPWQVPASASEVQIRTPRVNCPKVIICLDLSEEMSVPKLESFNGSRTNALNVSQKMVMFVRTKH
KIDKSHEFALVVNDDSAWLSGLTSDPRELCSCLYDLETASCSTFNLEGLFSLIQKTELPTVNTENVQTIP
PPYVVRTILVYSRPPCQPQFSLTEPMKMFQCPYFFFDIVYIHNGTTEEKEEDMSWKDMFAFMGSLDTKGA
SYKYEVALAGPALELHNCMAKLLAHLQRPCQTHSSYSLLEEDDEEAGEEEATV
```

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



ACCN: NM_026636

ORF Size: 1002 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_026636.1](#), [NP_080912.1](#)

RefSeq Size: 1592 bp

RefSeq ORF: 1002 bp

Locus ID: 68251


UniProt ID: [Q3UI43](#)

Cytogenetics: 8

MW: 36.8 kDa

Gene Summary: Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates. In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1. Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression. Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR204929