

Product datasheet for MR204987L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Abhd6 (NM_025341) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Abhd6 (NM 025341) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Abhd6

Synonyms: 0610041D24Rik; AA673485; AV065425

Mammalian Cell

viaitiitiailati CCII

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_025341 **ORF Size:** 1011 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR204987).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 025341.3, NP 079617.2

RefSeq Size: 2018 bp
RefSeq ORF: 1011 bp
Locus ID: 66082
UniProt ID: Q8R2Y0

Cytogenetics: 14 A1





Gene Summary:

Lipase that preferentially hydrolysis medium-chain saturated monoacylglycerols including 2-arachidonoylglycerol (PubMed:18096503, PubMed:20657592). Through 2-arachidonoylglycerol degradation may regulate endocannabinoid signaling pathways (PubMed:18096503, PubMed:20657592). Also has a lysophosphatidyl lipase activity with a preference for lysophosphatidylglycerol among other lysophospholipids (PubMed:24095738). Also able to degrade bis(monoacylglycero)phosphate (BMP) and constitutes the major enzyme for BMP catabolism (PubMed:26491015). BMP, also known as lysobisphosphatidic acid, is enriched in late endosomes and lysosomes and plays a key role in the formation of intraluminal vesicles and in lipid sorting (PubMed:26491015).[UniProtKB/Swiss-Prot Function]