

Product datasheet for MR205350L4

Ada (NM_007398) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ada (NM_007398) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Ada
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR205350).
Restriction Sites:	Sgfl-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_007398
ORF Size:	1059 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_007398.4 , NP_031424.1
RefSeq Size:	1701 bp
RefSeq ORF:	1059 bp
Locus ID:	11486
UniProt ID:	P03958
Cytogenetics:	2 H3
Gene Summary:	Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine (PubMed:9272950). Plays an important role in purine metabolism and in adenosine homeostasis (PubMed:9272950, PubMed:10720488). Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events (PubMed:11435465). Acts as a positive regulator of T-cell coactivation, by binding DPP4. Its interaction with DPP4 regulates lymphocyte-epithelial cell adhesion (By similarity). Enhances dendritic cell immunogenicity by affecting dendritic cell costimulatory molecule expression and cytokines and chemokines secretion (By similarity). Enhances CD4+ T-cell differentiation and proliferation (By similarity). Acts as a positive modulator of adenosine receptors ADORA1 and ADORA2A, by enhancing their ligand affinity via conformational change (By similarity). Stimulates plasminogen activation (By similarity). Plays a role in male fertility (By similarity). Plays a protective role in early postimplantation embryonic development (PubMed:9272950).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR205350L4