

## Product datasheet for MR205482L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cebpa (NM\_007678) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Cebpa (NM\_007678) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Cebpa

**Synonyms:** C/ebp; C/ebpalpha; CBF-A; Ceb; Cebp

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM\_007678

**ORF Size:** 1077 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(MR205482).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 007678.3, NP 031704.2

 RefSeq Size:
 2651 bp

 RefSeq ORF:
 1080 bp

 Locus ID:
 12606

 UniProt ID:
 P53566

 Cytogenetics:
 7 21.02 cM







## **Gene Summary:**

This intronless gene encodes a transcription factor that contains a basic leucine zipper (bZIP) domain and recognizes the CCAAT motif in the promoters of target genes. The encoded protein functions in homodimers and also heterodimers with CCAAT/enhancer-binding proteins beta and gamma. Activity of this protein can modulate the expression of genes involved in cell cycle regulation as well as in body weight homeostasis. The use of alternative in-frame non-AUG (CUG) and AUG start codons results in several protein isoforms with different lengths. Differential translation initiation is mediated by an out-of-frame, upstream open reading frame which is located between the CUG and the first AUG start codons. [provided by RefSeq, Sep 2014]