

Product datasheet for MR205511

Il1rap (NM_134103) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Il1rap (NM_134103) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Il1rap
Synonyms:	6430709H04Rik; AI255955; AV239853; IL-1RAcP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR205511 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGGACTTCTGTGGTATTTGATGAGTCTGTCCTTCTATGGGATCCTGCAGAGTCATGCTTCGGAGCGCT
GTGATGACTGGGGACTAGATACCATGCGACAAATCCAAGTGTGTTGAAGATGAGCCGGCTCGAATCAAGTG
CCCCCTTTTGAACACTTCTGAAGTACAACACAGCACTGCCATTCTCTGGCCTTACCCTGATCTGG
TACTGGACCAGGCAAGACCGGGACCTGGAGGAGCCATTAACCTCCGCCTCCAGAGAATCGCATCAGTA
AGGAGAAAAGATGTGCTCTGGTTCGGGCCACCCTCCTCAATGACACGGGCAATTACACCTGCATGTTGAG
GAACACAACCTTACTGCAGCAAAGTGCATTTCCCTGGAAGTTGTTTCAGAAGGACAGCTGTTTCAATTCT
GCCATGAGATTCCAGTGCAACAAGATGTATATTGAACATGGCATTTCATAAGATCAGATGCCAAATGTAG
ACGGATACTTTCCCTTCCAGTGTCAAACCATCGGTCACTTGGTATAAGGGTTGACTGAAATAGTGGACTT
TCATAATGTACTACCCGAGGGCATGAACTTGAGCTTTTTCATCCCTTGGTTTCAAATAACGGAAATTAC
ACATGTGTGGTTACATATCCTGAAAACGGACGTCTCTTTCACCTCACCAGGACTGTGACTGTAAAGGTGG
TGGGCTCACCAAAGGATGCATTGCCACCCAGATCTATTCTCAAATGACCGTGTGTCTATGAGAAAGA
ACCAGGAGAGGAACTGGTTATTCCTGCAAAGTCTATTTTCAGTTTCATTATGGACTCCCACAATGAGGTC
TGGTGGACCATTTGATGGAAGAAGCCTGATGACGTCACAGTCGACATCACTATTAATGAAAGTGAAGTT
ATTCTTCAACGGAAGATGAAACAAGGACTCAGATTTTTCAGCATCAAGAAAGTACCCCGGAGGATCTCAG
GCGCAACTATGTCTGTCTGCTGAAATACCAAAGGGGAAGCTGAGCAGGCTGCCAAGGTGAAACAGAAA
GGTAATGGATGCACCGAGCCGATGACTCTG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR205511 protein sequence
Red=Cloning site Green=Tags(s)

MGLLWYLMSLSFYGIQSHASERCDDWGLDTRQIQVFEDEPARIKCPLFEHFLKYNYSTAHSSGLTLIW
 YWTRQDRDLEEFNFRLENRISKEKDVLFWRPTLLNDTGNVTCMLRNTTYCSKVAFPLEVVQKDSFN
 AMRFPVHKMYIEHGIHKITCPNVDGYFPSSVKPSVTWYKGCETIVDFHNVLP EGMNLSFFIPLVSNNGNY
 TCVVYTPENGRFLHLTRVTVKVVGSPKDALPPQIYSPNDRVVYEKEPGEELVIPCKVYFSFIMDSHNEV
 WWTIDGKKPDDVTVDITINESVSYSTDETRTQILSIKKVTPEDLRNRYVCHARNTKGEAEQAQAVKQK
 GNGCTEPMTL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_134103

ORF Size: 1083 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_134103.2](#)

RefSeq Size: 2352 bp

RefSeq ORF: 1083 bp

Locus ID: 16180


UniProt ID: [Q61730](#)

Cytogenetics: 16 B2

MW: 41.6 kDa

Gene Summary: Coreceptor for IL1RL2 in the IL-36 signaling system. Coreceptor with IL1R1 in the IL-1 signaling system. Associates with IL1R1 bound to IL1B to form the high affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-kappa-B and other pathways. Signaling involves the recruitment of adapter molecules such as TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor subunits. Recruits TOLLIP to the signaling complex. Does not bind to interleukin-1 alone; binding of IL1RN to IL1R1, prevents its association with IL1R1 to form a signaling complex. The cellular response is modulated through a non-signaling association with the membrane IL1R2 decoy receptor. Secreted forms (isoforms 2 and 3) associate with secreted ligand-bound IL1R2 and increase the affinity of secreted IL1R2 for IL1B; this complex formation may be the dominant mechanism for neutralization of IL1B by secreted/soluble receptors. Coreceptor for IL1RL1 in the IL-33 signaling system. Can bidirectionally induce pre- and postsynaptic differentiation of neurons by trans-synaptically binding to PTPRD (PubMed:25908590). May play a role in IL1B-mediated costimulation of IFNG production from T-helper 1 (Th1) cells (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR205511