

## Product datasheet for MR205751

### Kcnj1 (NM\_019659) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Kcnj1 (NM_019659) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Kcnj1
Synonyms:	Kir1.1; ROMK; Romk2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR205751 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGTTCAAACATCTTCGAAGATGGTTTGTCACTCACATATTTGGGCGTTCTCGGCAACGAGCAAGTTGG  
TCTCCAAAGATGGAAGGTGTAACATCGAGTTTGGCAATGTAGATGCACAGTCGAGGTTTATATTCTTTGT  
GGATATCTGGACAACTGACTTGACCTGAAATGGAGGTACAAAATGACCGTGTTTCATCACAGCCTTCTTG  
GGGAGTTGGTTTCTCTTTGGTCTCCTGTGGTATGTCGTAGCCTATGTTTCATAAGGATCTCCAGAGTTCT  
ACCCACCTGACAACCGTACTCCTTGTGTGGAGAACATTAATGGCATGACATCAGCCTTTCTGTTTTCTCT  
AGAGACCCAAGTGACCATAGGTTACGGATTCAGGTTTGTGACAGAACAGTGTGCCACTGCCATTTTTCTG  
CTTATCTTCCAGTCTATTCTTGGAGTGATCATCAATTCTTTCATGTGTGGTGCCATATTAGCCAAGATCT  
CTAGACCCAAAAACGTGCAAAGACATTACATTCAGCAAGAAATGCGGTGATCAGCAAACGTGGGGGAA  
GCTCTGTCTCCTCATCCGAGTAGCAAACTTTAGGAAAAGCCTTCTGATTGGCAGTCACATATATGGTAAG  
CTTCTGAAGACTACCATCACACCTGAAGGAGAGACCATTATTTGGATCAGACCAATATAAACTTTGTAG  
TTGATGCTGGCAATGAAAATTTGTTCTTCAATTTCCCACTGACAATCTACCACATTATTGACCACAACAG  
CCCTTTCTTCCACATGGCGGCAGAACTCTTCCCAACAGGACTTCGAGTTGGTTGTCTTTTAGATGCC  
ACAGTAGAATCCACAGTGCAACCTGCCAAGTCCGCACATACATCCCAGAAGAGGTGCTTTGGGTT  
ACCGTTTTGTCCCATCGTATCCAAGACCAAGGAAGGAAATACCGAGTGGATTTCCATAACTTTGGTAA  
GACGGTGAAGTGAGACCCCTCATTGTGCCATGTGCCTATAATGAGAAAGATGCCAGGGCCAGGATG  
AAGAGAGGCTATGACAACCCTAACCTTTGTCTTGTGAGAAGTTGATGAAACAGACGACACCCAAATG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >MR205751 protein sequence  
Red=Cloning site Green=Tags(s)

MFKHLRRWFVTHIFGRSRQRARLVSKDGRCNIEFGNVDAQSRFIFVVDIWTTLVLDLKWRYKMTVFITAFL  
 GSWFLFGLLWYVAVYVHKDLPEFYPPDNRTPCVENINGMTSAFLFSLAQVTIGYGFRFVTEQCATAIFL  
 LIFQSIILGVIINSFMCGAILAKISRPKKRAKTITFSKNAVISKRGGKLCLLIRVANLRKSLIGSHIYGK  
 LLKTTITPEGETIILDQTNINFVVDAGNENLFFISPLTIYHIIDHNSPFFHMAAETLSQQDFELVVFLDG  
 TVESTSATCQVRTSYIPEEVLWGYRFVPIVSKTKEGKYRVDFHNFGKTVEVETPHCAMCLYNEKDARARM  
 KRGYDNPNFVLSEVDETDQTM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:**

SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF

**ACCN:** NM\_019659

**ORF Size:** 1119 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_019659.3](#)

**RefSeq Size:** 3073 bp

**RefSeq ORF:** 1119 bp

**Locus ID:** 56379


**UniProt ID:** [O88335](#)

**Cytogenetics:** 9 A4

**MW:** 42.8 kDa

**Gene Summary:** In the kidney, probably plays a major role in potassium homeostasis. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. This channel is activated by internal ATP and can be blocked by external barium (By similarity).[UniProtKB/Swiss-Prot Function]

**Product images:**



Circular map for MR205751