

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR205897L3V

Prmt6 (NM_178891) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Prmt6 (NM_178891) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Prmt6
Synonyms:	AW124876; BB233495; Hrmt1l6
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_178891
ORF Size:	1134 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR205897).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 178891.5, NP 849222.3</u>
RefSeq Size:	2475 bp
RefSeq ORF:	1137 bp
Locus ID:	99890
UniProt ID:	<u>Q6NZB1</u>
Cytogenetics:	3 F3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Arginine methyltransferase that can catalyze the formation of both omega-N Gene Summary: monomethylarginine (MMA) and asymmetrical dimethylarginine (aDMA), with a strong preference for the formation of aDMA (PubMed:22904064, PubMed:26070566). Preferentially methylates arginyl residues present in a glycine and arginine-rich domain and displays preference for monomethylated substrates (By similarity). Specifically mediates the asymmetric dimethylation of histone H3 'Arg-2' to form H3R2me2a (By similarity). H3R2me2a represents a specific tag for epigenetic transcriptional repression and is mutually exclusive with methylation on histone H3 'Lys-4' (H3K4me2 and H3K4me3) (By similarity). Acts as a transcriptional repressor of various genes such as HOXA2, THBS1 and TP53 (PubMed:22904064). Repression of TP53 blocks cellular senescence (PubMed:22904064). Also methylates histone H2A and H4 'Arg-3' (H2AR3me and H4R3me, respectively). Acts as a regulator of DNA base excision during DNA repair by mediating the methylation of DNA polymerase beta (POLB), leading to the stimulation of its polymerase activity by enhancing DNA binding and processivity. Methylates HMGA1. Regulates alternative splicing events. Acts as a transcriptional coactivator of a number of steroid hormone receptors including ESR1, ESR2, PGR and NR3C1. Promotes fasting-induced transcriptional activation of the gluconeogenic program through methylation of the CRTC2 transcription coactivator (PubMed:24570487). Methylates GPS2, protecting GPS2 from ubiquitination and degradation (PubMed:26070566).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US