

Product datasheet for MR206198

Vrk1 (NM_001029844) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Vrk1 (NM_001029844) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Vrk1
Synonyms:	51PK
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR206198 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGCCCGGTGTAAGAGCAGCTCAGGCTGGAAGACCCGGACCTGCGAAGAGGCGCCTCGCAGAGCAGTTTG
CCGCTGGAGAGGTCCTAACCGACATGTCTAGGAAGGAGTGAAACTAGGATTGCCATTGGCCAAGGTGG
CTTTGGCTGCATCTATCTGGCGGACACAAATCTTCCAAACCGGTTGGCAGTGACGCGCCCTGTGTGTG
AAAGTGAACCCAGTGACAATGGACCTCTTTTCACGGAATTAAGTTCTACCAGAGGGCTGCTAAACCAG
AGCAAATTCAGAAATGGATTTCGTACACATAAATTGAAGTACCTTGGTGTTCCTAAGTATTGGGGATCTGG
TCTACATGATAAAAATGAAAAAGTTACAGTTTTATGATAATGGACCGCTTTGGGAGTGACCTTCAGAAA
ATATATGAAGCAAATGCCAAAAGTTTTCTCGGAAAACCTGATTGCAGCTAAGCTTAAGAATTCTGGATA
TCCTGGAGTACATCCACGAGCATGAGTACGTGCACGGGGACATCAAGGCCTCCAACCTGCTCCTGAGTCA
CAAGAACCCTGACCAGGTATATTTGGTAGACTATGGCCTTGCTTATCGGTACTGCCAGATGGAGTTCAT
AAAGAGTACAAGGAAGATCCCAAAGGTGCCATGACGGCACCCCTGGAGTTCACCAGCATCGACGCTCACA
AAGCGTGGCCCCATCAAGACGTGGTGATTTGAAATACTTGGTTATTGCATGATCCAGTGGCTCAGCGG
CTGTCTTCCTTGGGAAGATAACTTGAAGATCCTAACTACGTTAGGGATTCCAAAATTAGATACAGAGAC
AACGTCGCAGCTTTGATGGAGAAATGCTTTCCTGAGAAAAATAAGCCAGGTGAGATCGCTAAGTACATGG
AGTCTGTGAAACTACTGGAATACACCGAAAAACCTCTCTATCAAAACCTACGTGATATCCTTTTACAAGG
ACTAAAAGCTATAGGAAGTAAAGACGACGGCAAACCTGGATTTTAGTGCTGTGGAGAACGGAAGTGTGAAG
ACAAGACCAGCCTCAAAGAAGCGGAAGAAAGAAGCAGAAGAAAGCGCGGTGTGCGCTGTGGAGGACATGG
AGTGCTCAGACACACAGGTGCAAGAGGCCCGCACAGACCCGTTACGAACCAGAAAGAAAGCCAGAAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR206198 protein sequence
Red=Cloning site Green=Tags(s)

MPRVKAAQAGRPGPAKRRLAEQFAAGEVLTDMSRKEWKLGLPIGQGGFGCIYLADTNSSKPVGSDAPCVV
 KVEPSDNGPLFTELKQYRAAKPEQIQKWIRTHKLYLGVPKYWGSLHDKNGKSYRFMIMDRFGSDLQK
 IYEANAKRFSRKTVLQLSLRILDILEYIHEHEYVHGDIKASNLLL SHKNPDQVYLVDYGLAYRYCPDGVH
 KEYKEDPKRCHDGTLEFTSIDAHKGVAPSRRGDLEILGYCMIQWLSGCLPWEDNLKDPNYVRDSKIRYRD
 NVAALMEKCFPEKNKPGEIAKYMESVKLLEYTEKPLYQNLRDILLQGLKAI GSKDDGKLD FSAVENG SVK
 TRPASKRRKKEAESAVCAVEDMECSDTQVQEAQAQTRSRTTRKKAQK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001029844

ORF Size: 1191 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001029844.1](#), [NP_001025015.1](#)

RefSeq Size: 3664 bp

RefSeq ORF: 1191 bp

Locus ID: 22367


UniProt ID: [Q80X41](#)

Cytogenetics: 12 E- F1

MW: 45 kDa

Gene Summary: Serine/threonine kinase involved in Golgi disassembly during the cell cycle: following phosphorylation by PLK3 during mitosis, required to induce Golgi fragmentation. Acts by mediating phosphorylation of downstream target protein. Phosphorylates 'Thr-18' of p53/TP53 and may thereby prevent the interaction between p53/TP53 and MDM2. Phosphorylates casein and histone H3. Phosphorylates BANF1: disrupts its ability to bind DNA, reduces its binding to LEM domain-containing proteins and causes its relocalization from the nucleus to the cytoplasm. Phosphorylates ATF2 which activates its transcriptional activity (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR206198