

## Product datasheet for MR206384L4

### Snx6 (NM\_026998) Mouse Tagged Lenti ORF Clone

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Expression Plasmids  |
| Product Name:             | Snx6 (NM_026998) Mouse Tagged Lenti ORF Clone                  |
| Tag:                      | mGFP   |
| Symbol:                   | Snx6   |
| Synonyms:                 | 2010006G21Rik; 2610032J07Rik; 2810425K19Rik; AU018928; C85963  |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)                              |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)                                     |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR206384). |
| Restriction Sites:        | SgfI-MluI  |
| Cloning Scheme:           |  |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |           |
|-----------|-----------|
| ACCN:     | NM_026998 |
| ORF Size: | 1218 bp   |



[View online »](#)

|                               |   |
|-------------------------------|---|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>  |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>   |
| <b>RefSeq:</b>                | <a href="#">NM_026998.2</a> , <a href="#">NP_081274.2</a>   |
| <b>RefSeq Size:</b>           | 1880 bp   |
| <b>RefSeq ORF:</b>            | 1221 bp   |
| <b>Locus ID:</b>              | 72183   |
| <b>UniProt ID:</b>            | <a href="#">Q6P8X1</a>  |
| <b>Cytogenetics:</b>          | 12 C1   |
| <b>Gene Summary:</b>          | Involved in several stages of intracellular trafficking. Interacts with membranes phosphatidylinositol 3,4-bisphosphate and/or phosphatidylinositol 4,5-bisphosphate (Probable). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC). Does not have in vitro vesicle-to-membrane remodeling activity (By similarity). Involved in retrograde endosome-to-TGN transport of lysosomal enzyme receptor IGF2R. May function as link between transport vesicles and dynactin. Negatively regulates retrograde transport of BACE1 from the cell surface to the trans-Golgi network. Involved in E-cadherin sorting and degradation; inhibits PIP5K1C-mediated E-cadherin degradation (By similarity). In association with GIT1 involved in EGFR degradation (PubMed:18523162). Promotes lysosomal degradation of CDKN1B (PubMed:20228253). May contribute to transcription regulation (By similarity).[UniProtKB/Swiss-Prot Function] |

## Product images:



Circular map for MR206384L4