

Product datasheet for MR206404L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Adrm1 (NM_019822) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Adrm1 (NM_019822) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Adrm1

Synonyms: 1110063P18Rik; 2510006J17Rik; AA408205; ARM-1; Arm1; AU043535; Gp110; Rpn13

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_019822 **ORF Size:** 1224 bp

ORF Nucleotide

•

Sequence:

The ORF insert of this clone is exactly the same as(MR206404).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 019822.3, NP 062796.2

RefSeq Size: 1429 bp
RefSeq ORF: 1224 bp
Locus ID: 56436
UniProt ID: Q9JKV1

Cytogenetics: 2 H4







Gene Summary:

Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. Within the complex, functions as a proteasomal ubiquitin receptor. Engages and thus activates 19S-associated deubiquitinases UCHL5 and PSMD14 during protein degradation. UCHL5 reversibly associate with the 19S regulatory particle whereas PSMD14 is an intrinsic subunit of the proteasome lid subcomplex.[UniProtKB/Swiss-Prot Function]