

Product datasheet for MR206531L3V

Yy1 (NM_009537) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles Product Name: Yy1 (NM 009537) Mouse Tagged ORF Clone Lentiviral Particle Symbol: Yv1 AW488674; NF-E1 Synonyms: **Mammalian Cell** Puromycin Selection: Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092) Tag: Myc-DDK NM 009537 ACCN: ORF Size: 1242 bp The ORF insert of this clone is exactly the same as(MR206531). **ORF** Nucleotide Sequence: **OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info **OTI** Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. **RefSeq:** NM 009537.3 **RefSeq Size:** 2324 bp **RefSeq ORF:** 1245 bp Locus ID: 22632 **UniProt ID:** Q00899 Cytogenetics: 12 59.58 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE Yy1 (NM_009537) Mouse Tagged ORF Clone Lentiviral Particle – MR206531L3V

Multifunctional transcription factor that exhibits positive and negative control on a large Gene Summary: number of cellular and viral genes by binding to sites overlapping the transcription start site. Binds to the consensus sequence 5'-CCGCCATNTT-3'; some genes have been shown to contain a longer binding motif allowing enhanced binding; the initial CG dinucleotide can be methylated greatly reducing the binding affinity. The effect on transcription regulation is depending upon the context in which it binds and diverse mechanisms of action include direct activation or repression, indirect activation or repression via cofactor recruitment, or activation or repression by disruption of binding sites or conformational DNA changes. Its activity is regulated by transcription factors and cytoplasmic proteins that have been shown to abrogate or completely inhibit YY1-mediated activation or repression. Binds to the upstream conserved region (UCR) (5'-CGCCATTTT-3') of Moloney murine leukemia virus (MuLV). Acts synergistically with the SMAD1 and SMAD4 in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression (PubMed:15329343). Binds to SMAD binding elements (SBEs) (5'-GTCT/AGAC-3') within BMP response element (BMPRE) of cardiac activating regions (PubMed:15329343). Proposed to recruit the PRC2/EED-EZH2 complex to target genes that are transcriptional repressed. Involved in DNA repair. In vitro, binds to DNA recombination intermediate structures (Holliday junctions). Involved in spermatogenesis and may play a role in meiotic DNA double-strand break repair. Plays a role in regulating enhancer activation (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US