

Product datasheet for MR206564

Dclre1b (NM_001025312) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Dclre1b (NM_001025312) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Dclre1b
Synonyms:	AI452214; Apollo; mSNM1B; SNMIB
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR206564 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCTGAAGGAACCTGCTCTGATACTGGGAAACAGATTCATACTTTGTATCTAGACAACACTAATTGCA
ATCCAGCCTGGTTCTTCTTCCCGACAAGAGGCTACTCAGCAGATTGTCCAGCTAATCCGACAGTTCCC
ACAACACAACATAAAAAATTGGACTCTATAGTCTAGGAAAAGAATCACTGCTGGAGCAACTAGCCCTTGAG
TTTCGGACCTGGGTGGTATTAAGTCTCAACGCCTGGAGTTGGTACAGTTGCTGGGCCTGGCAGACGTGT
TCACAGTTGAGGAAGAAGCCGGGCGCATCCATGCTGTGGACCATACGGAGATCTGCCACTCTGCCATGCT
TCAGTGAACAGAGTCACCCTACCATTGCTATTTCCCAAGCCGAAAGTGGCAGCCCTCACCCC
AGCATCTACACCGTCCCCTACTCTGACCATTCTTACTCCGAGCTCCGTGCTTTTGTGGCAGCGCTGA
GGCCTTGTGAGTGGTGGCCATCGTCCATCAAAAGCCTTGTGGAGAGTTCTTTCAGGACAGCTTGAGTCC
TAGGCTCGCTATGCCTCTGATTCCACACTCTGTGCAGCAGTACATGAGTTCTCTCTAGGAAAACAAAT
GTGCTTTGGCAGTTAGAAAGGAGGCTCAAGAGGCCAAGAAGTCAAGGTTGTGTTTGAATCCCCTGAGG
AGAAAGCTAATCAGGTTAAAGTTGACAGAGACTCAAAGAAGCAGAAAAGGAAAACCTCTCTCCCTGGGC
TGGCACCTTGAGAGGCTTTGCCCATCCTCTGCAGGCCAGGAAGCAGTTGTTCCAGACTTCTGCAGG
AAAGAAGCTGATGAGCCAGTCTCTTTTGTGACTCCAACAAGTGGCAGCTGTGTTGACTGCCCCACTGG
AATTCTCAGTGCAATTACAACCTATAGATGAGTTTCTCTTCCAGAAACCAGGGAGAAAATTGGCTTAGA
GTCCCCATTGCTGCAAGGGGAGACAGTGGCTCACAGCGAGAGGGAACCAAAGTACTGCGTAGGCTGT
GGTTCTCCCCGGCTCACATTAGCAGAGCTGTTCTCTAACTCTGAGTCCAGGGGCTGGCGCTAAAAAT
ACCTTCTCACTCCAGTGCATTTCTCCAAGCAGGCTTCTCTTCCAGGAACCTCGACAAGCAAGTGGAAAA
ACACCAGAGAGTACAGCGTAGCAGTCTGCAGTCTGAGCCCGGTGGATGTTGGC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online >](#)

Protein Sequence: >MR206564 protein sequence
Red=Cloning site Green=Tags(s)

MLKEPALILGKQIHTLYLDNTNCNPALVLP SRQEATQQIVQLIRQFPQHNIKIGLYSLGKESLLEQLALE
 FRTWVVLSPQRLELVQLGLADVFTVEEEAGRIHAVDHTEICH SAMLQWNQSHPTIAIFPTSRKVRSPHP
 SIYTVPYSDHSSYSELRAFVAALRPCQVVPVIVHQKPCGEFFQDSLSPRLAMPLIPHSVQQYMSSSSRKTN
 VLWQLERRLKRPR TQGVVFE SPEEKANQVKVDRDSKHKHKKENLSPWAGHLERLCPHP LQARKQLFPDFCR
 KERDEPVLFCDSNKMATVLTAPLEFSVQLQPIDEF LFPETREKIGLESPLL SRGDSGSPARGNQSDCVGC
 GSPPAHISR AVPLTPESRGLALKYLLTPVHFLQAGFSSRNFDKQVEKHQRVQRSSPAVLSPVDVG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001025312

ORF Size: 1248 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001025312.1](#), [NP_001020483.1](#)

RefSeq Size: 4143 bp

RefSeq ORF: 1248 bp

Locus ID: 140917


UniProt ID: [Q8C7W7](#)

Cytogenetics: 3 F2.2

MW: 46.9 kDa

Gene Summary: 5'-3' exonuclease that plays a central role in telomere maintenance and protection during S-phase. Participates in the protection of telomeres against non-homologous end-joining (NHEJ)-mediated repair, thereby ensuring that telomeres do not fuse. Plays a key role in telomeric loop (T loop) formation by being recruited by TERF2 at the leading end telomeres and by processing leading-end telomeres immediately after their replication via its exonuclease activity: generates 3' single-stranded overhang at the leading end telomeres avoiding blunt leading-end telomeres that are vulnerable to end-joining reactions and expose the telomere end in a manner that activates the DNA repair pathways. Together with TERF2, required to protect telomeres from replicative damage during replication by controlling the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B) needed for telomere replication during fork passage and prevent aberrant telomere topology. Also involved in response to DNA damage: plays a role in response to DNA interstrand cross-links (ICLs) by facilitating double-strand break formation. In case of spindle stress, involved in prophase checkpoint. [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR206564