

## Product datasheet for MR206648L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Kcnj16 (NM\_010604) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Kcnj16 (NM 010604) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Kcnj16

**Synonyms:** 6430410F18Rik; AI132396; Kir5.1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_010604 **ORF Size:** 1260 bp

**ORF Nucleotide** 

The ODE

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(MR206648).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 010604.3

 RefSeq Size:
 3623 bp

 RefSeq ORF:
 1260 bp

 Locus ID:
 16517

 UniProt ID:
 Q9Z307

Cytogenetics: 11 75.01 cM







## **Gene Summary:**

Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. KCNJ16 may be involved in the regulation of fluid and pH balance. In the kidney, together with KCNJ10, mediates basolateral K(+) recycling in distal tubules; this process is critical for Na(+) reabsorption at the tubules.[UniProtKB/Swiss-Prot Function]