

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR206727L3V

Cops3 (NM_011991) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cops3 (NM_011991) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Cops3
Synonyms:	Csn3; Sgn3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_011991
ORF Size:	1272 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR206727).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 011991.1, NP 036121.1</u>
RefSeq Size:	1583 bp
RefSeq ORF:	1272 bp
Locus ID:	26572
UniProt ID:	<u>088543</u>
Cytogenetics:	11 B1.3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Component of the COP9 signalosome complex (CSN), a complex involved in various cellular
and developmental processes. The CSN complex is an essential regulator of the ubiquitin
(Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type
E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as
SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN,
IkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD
kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects
degradation by the Ubl system, respectively. Essential to maintain the survival of epiblast cells
and thus the development of the postimplantation embryo.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US