

Product datasheet for MR206727L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cops3 (NM_011991) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cops3 (NM_011991) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Cops3

Synonyms: Csn3; Sgn3

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_011991 **ORF Size:** 1272 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR206727).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 011991.1, NP 036121.1

 RefSeq Size:
 1583 bp

 RefSeq ORF:
 1272 bp

 Locus ID:
 26572

 UniProt ID:
 088543

 Cytogenetics:
 11 B1.3







Gene Summary:

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the UbI ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, lkappaBalpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the UbI system, respectively. Essential to maintain the survival of epiblast cells and thus the development of the postimplantation embryo.[UniProtKB/Swiss-Prot Function]