

Product datasheet for MR206733L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Acp2 (NM_007387) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Acp2 (NM 007387) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Acp2

Synonyms: Acp; Acp-2; LAP

Mammalian Cell Pe

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_007387 **ORF Size:** 1272 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR206733).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 007387.1

 RefSeq Size:
 4669 bp

 RefSeq ORF:
 1272 bp

 Locus ID:
 11432

 UniProt ID:
 P24638

 Cytogenetics:
 2 50.54 cM







Gene Summary:

The protein encoded by this gene belongs to the histidine acid phosphatase family, which hydrolyze orthophosphoric monoesters to alcohol and phosphate. This protein is localized to the lysosomal membrane, and is chemically and genetically distinct from the red cell acid phosphatase. Mice lacking this gene showed multiple defects, including bone structure alterations, lysosomal storage defects, and an increased tendency towards seizures. An enzymatically-inactive allele of this gene showed severe growth retardation, hair-follicle abnormalities, and an ataxia-like phenotype. Two isoforms are predicted to be produced from the same mRNA by the use of alternative in-frame translation termination codons via a stop codon readthrough mechanism. [provided by RefSeq, Oct 2017]