

Product datasheet for **MR206782L4V**

Aadat (NM_011834) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Aadat (NM_011834) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Aadat
Synonyms:	Aadt; A1875679; Kat2; KATII; Kyat2; mKat-2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_011834
ORF Size:	1278 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR206782).
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_011834.2 , NP_035964.1
RefSeq Size:	1780 bp
RefSeq ORF:	1278 bp



[View online »](#)

Locus ID: 23923

UniProt ID: [Q9WVM8](#)

Cytogenetics: 8 30.85 cM

Gene Summary: Transaminase with broad substrate specificity. Has transaminase activity towards amino adipate, kynurenine, methionine and glutamate. Shows activity also towards tryptophan, aspartate and hydroxykynurenine. Accepts a variety of oxo-acids as amino-group acceptors, with a preference for 2-oxoglutarate, 2-oxocaproic acid, phenylpyruvate and alpha-oxo-gamma-methyl butyric acid. Can also use glyoxylate as amino-group acceptor (in vitro) (By similarity).[UniProtKB/Swiss-Prot Function]