

Product datasheet for **MR206809**

Stk3 (BC037440) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Stk3 (BC037440) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Stk3
Synonyms:	Mess1, Mst3, MST
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>MR206809 representing BC037440 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGAGCAGCCGCCGGCTCCAAGAGTCCATATGTTGTGAAGTACTATGGCAGTTACTTTAAGAACACAG
ACCTCTGGATTGTTATGGAGTACTGTGGAGCGGGTCCGTTTCAGACATAATTAGATTGCGAAACAAGAC
ATTAACAGAAGATGAAATTGCAACTATTCTAAAATCCACATTGAAAGGATTAGAATATTTGCATTTTATG
AGGAAAATACACAGAGATATAAAGCCGGGAATATTCTCCTCAATACAGAAGGACATGCAAAGCTTGCGAG
ATTTTGGAGTGGCTGGCCAGTTAACAGATACAATGGCAAAACGCAACACTGTAATAGGAACCCCATTTTG
GATGGCTCCTGAGGTAATTCAAGAAATAGGTTACAACCTGTGTGGCTGACATCTGGTCCCTTGGCATTACT
TCTATAGAAATGGCAGAAGGAAACCTCCTTATGCTGATATACATCCGATGAGGGCTATTTTTATGATCC
CTACAAACCCACCACCAACATTCAGGAAACCTGAACTTTGGTCTGATGACTTCACCGATTTTGTGAAGAA
GTGCTTGGTGAAGAGTCTGAGCAGAGAGCCACTGCGACACAGCTGTTACAGCATCCTTTTATCAAGAAT
GCAAAACCCGTGTCGATATTAAGAGACCTGATCGCAGAAGCCATGGAAATCAAAGCCAAACGGCATGAAG
AGCAGCAGCGGGAGCTAGAGGAGGAGGAAAGAAACTCGGATGAAGATGAGCTGGATTACACACTATGGT
GAAGACGAGTTTCAGAAAGCGTGGGCACAATGCGGGCCACCAGCAGCATGAGTGAAGGAGCCAGACGATG
ATTGAACATAACAGCACCATGTTAGAGTCGGACCTGGGGACCATGTTTATAAACAGTGAAGAAGAGGAGG
AAGAGGAAGAGG
TTCTTTTATGACTACTTTGATAAGCAGGACTTCAAGAACAAGAGTCATGAAAATTTGTGATCAGAGCATG
CGTGAGCCAGGCCCTATGTCCAACAGTGTTTTTCTGACAACCTGGAGAGTTCTCAAGATGGAGACTTTG
ATTTCTTAAAAATCTAAGTTTAGAAGAACTACAGATGCGGCTAAAAGCACTAGACCCCATGATGGAACG
AGAAATAGAAGAACTGCATCAAAGATACAGTGCAAAAAGACAACCTATCCTGGATGCCATGGATGCGAAG
AAGAGGAGGCAGCAGAATTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR206809 representing BC037440
Red=Cloning site Green=Tags(s)



MEQPPASKSPYVVKYGSYFKNTDLWIVMEYCGAGSVSDIIRLRNKLTLEDEIATILKSTLKGLEYLHFM
 RKIHRDIKAGNILLNTEGHAKLADFGVAGQLTDTMAKRNTVIGTPFWMAPEVIQEI GYNCVADIWSLGIT
 SIEMAEKGPPYADIHPMRAIFMIPTNPPPTFRKPELWSDDFDFVKKCLKVSPEQRATATQLLQHPFIKN
 AKPVSILRDLIAEAMEIKAKRHEEQQRELEEEEEENSDDELDSHTMVKTSSES SVGTM RATSTMSEGAQTM
 IEHNSTMLES DLGTMVINSEEEEEEEEEEDGTMKRNATSPQVQRPSFMDYFDKQDFKNKSHENCDSM
 REPGPMSNSVFPDNWRVPQDGFDFLKNLSLEELQMRLKALDPMEREIEELHQRYSAKRQPILDAMDAK
 KRRQQNF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:
Cloning Scheme:


SgfI-MluI

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: BC037440

ORF Size:	1281 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC037440.1
RefSeq Size:	2495 bp
RefSeq ORF:	1283 bp
Locus ID:	56274
Cytogenetics:	15 B3.1
MW:	91.4 kDa

Gene Summary:

Stress-activated, pro-apoptotic kinase which, following caspase-cleavage, enters the nucleus and induces chromatin condensation followed by internucleosomal DNA fragmentation. Key component of the Hippo signaling pathway which plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Phosphorylation of YAP1 by LATS2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. STK3/MST2 and STK4/MST1 are required to repress proliferation of mature hepatocytes, to prevent activation of facultative adult liver stem cells (oval cells), and to inhibit tumor formation. Phosphorylates NKX2-1. Phosphorylates NEK2 and plays a role in centrosome disjunction by regulating the localization of NEK2 to centrosomes, and its ability to phosphorylate CROCC and CEP250. In conjunction with SAV1, activates the transcriptional activity of ESR1 through the modulation of its phosphorylation. Positively regulates RAF1 activation via suppression of the inhibitory phosphorylation of RAF1 on 'Ser-259'. Phosphorylates MOBKL1A and RASSF2. Phosphorylates MOBKL1B on 'Thr-74'. Acts cooperatively with MOBKL1B to activate STK38 (By similarity).[UniProtKB/Swiss-Prot Function]