

Product datasheet for MR206951L3

Sugct (NM_138654) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Sugct (NM_138654) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Sugct
Synonyms:	5033411D12Rik; D17907
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR206951).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_138654
ORF Size:	1311 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_138654.2 , NP_619595.2
RefSeq Size:	1727 bp
RefSeq ORF:	1311 bp
Locus ID:	192136
UniProt ID:	Q7TNE1
Cytogenetics:	13 A2
Gene Summary:	Catalyzes the succinyl-CoA-dependent conversion of glutarate to glutaryl-CoA. Can use different dicarboxylic acids as CoA acceptors, the preferred ones are glutarate, succinate, adipate, and 3-hydroxymethylglutarate (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR206951L3