

# **Product datasheet for MR206962L3V**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Tfap2a (NM\_011547) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Tfap2a (NM\_011547) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Tfap2a

Synonyms: A; AP; AP-2; Ap-2 (a); Ap2; AP2alpha; Ap2tf; Tcfa; Tcfap2a

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_011547

ORF Size: 1311 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR206962).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 011547.2, NP 035677.2

 RefSeq Size:
 2863 bp

 RefSeq ORF:
 1320 bp

 Locus ID:
 21418

 UniProt ID:
 P34056

Cytogenetics: 13 20.01 cM







#### **Gene Summary:**

This gene is a member of the activator protein 2 (AP-2) transcription factor family. The protein encoded by this gene can act as both an activator and repressor of gene transcription, and plays an important role in early embryogenesis, specifically in cranial development. This protein forms both homodimers and heterodimers, and binds to a GC-rich consensus sequence found in some promoters and enhancers. Disruption of this gene causes perinatal death, with neural tube, craniofacial, and limb mesenchyme defects. Alternative splicing results in multiple transcript variants that encode multiple protein isoforms. [provided by RefSeq, Sep 2014]