

Product datasheet for MR206965L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Vps4a (NM_126165) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Vps4a (NM_126165) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Vps4a

Synonyms: 4930589C15Rik; Al325971; AW553189

Mammalian Cell

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 126165

Tag: Myc-DDK

ORF Size: 1311 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR206965).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 126165.1, NP 569053.1

 RefSeq Size:
 2089 bp

 RefSeq ORF:
 1314 bp

 Locus ID:
 116733

 UniProt ID:
 Q8VEJ9

Cytogenetics: 8 D3





Gene Summary:

Involved in late steps of the endosomal multivesicular bodies (MVB) pathway. Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly, possibly in combination with membrane fission. Redistributes the ESCRT-III components to the cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. In conjunction with the ESCRT machinery also appears to function in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. Involved in cytokinesis: retained at the midbody by ZFYVE19/ANCHR and CHMP4C until abscission checkpoint signaling is terminated at late cytokinesis. It is then released following dephosphorylation of CHMP4C, leading to abscission. VPS4A/B are required for the exosomal release of SDCBP, CD63 and syndecan (By similarity).[UniProtKB/Swiss-Prot Function]